

TAF MEK melanoma - Management of patients with BRAF-positive melanoma - HCP

[Prescribing information](#)

Image



Image



## Management of patients with ***BRAF***-positive melanoma

TAFINLAR® (dabrafenib) in combination with MEKINIST® (trametinib) is indicated in adult patients with unresectable or metastatic melanoma with a *BRAF* V600 mutation.<sup>1,2</sup>

TAFINLAR in combination with MEKINIST is indicated for the adjuvant treatment of adult patients with Stage III melanoma with a *BRAF* V600 mutation, following complete resection.<sup>1,2</sup>

**For the full safety profile, please refer to the Summary of Product Characteristics (SmPC) for [TAFINLAR](#) and [MEKINIST](#).**

**Adverse event reporting: Details of how to report adverse events are available at the bottom of the page. Please refer to the respective SmPC for all licensed indications.**

## **Helping patients on their TAFINLAR + MEKINIST treatment journey**

**TAFINLAR + MEKINIST offers oral dosing. TAFINLAR is taken twice daily, whilst MEKINIST is taken once a day, with the morning or evening dose of TAFINLAR<sup>1,2</sup>**

TAFINLAR + MEKINIST offers oral dosing. It is the only oral option in the adjuvant setting and offers a low pill burden to patients in the metastatic setting.<sup>1-4</sup> Dosing is the same for both adjuvant and metastatic treatment settings.<sup>1,2</sup>

Please select an option below to learn more about dosing and administration, and how to manage adverse events with dose modifications.

*BRAF* V600, mutation of the *BRAF* gene at valine (V) 600; SmPC, summary of product characteristics.

### References

1. TAFINLAR (dabrafenib) Summary of Product Characteristics.
2. MEKINIST (trametinib) Summary of Product Characteristics.
3. BRAFTOVI (encorafenib) Summary of Product Characteristics.
4. MEKTOVI (binimetinib) Summary of Product Characteristics.

## **TAFINLAR + MEKINIST dosing and administration**

[Show more](#)

## **TAFINLAR + MEKINIST dose modifications**

[Show more](#)

UK | April 2025 | FA-11392338

Adverse events should be reported. Reporting forms and information can be found at [www.mhra.gov.uk/yellowcard](http://www.mhra.gov.uk/yellowcard). Adverse events should also be reported to Novartis online through the pharmacovigilance intake (PVI) tool at [www.novartis.com/report](http://www.novartis.com/report), or alternatively email [medinfo.uk@novartis.com](mailto:medinfo.uk@novartis.com) or call 01276 698370.

---

**Source URL:**

<https://www.pro.novartis.com/uk-en/medicines/oncology/tafinlar-mekinist/melanoma/management>