

Ilaris - Safety profile - HCP

[Prescribing information](#)

Image



Image



**ILARIS®**  
(canakinumab)  
150 mg subcutaneous injection

## **ILARIS® (canakinumab) safety profile**

### **Indications<sup>1</sup>**

#### Periodic fever syndromes

ILARIS is indicated for the treatment of the following autoinflammatory periodic fever syndromes in adults, adolescents and children aged 2 years and older:

- Cryopyrin-associated periodic syndromes (CAPS), including:
  - Muckle-Wells syndrome (MWS)

- Neonatal-onset multisystem inflammatory disease (NOMID)/chronic infantile neurological, cutaneous, articular syndrome (CINCA)
- Severe forms of familial cold autoinflammatory syndrome (FCAS)/familial cold urticaria (FCU) presenting with signs and symptoms beyond cold-induced urticarial skin rash
- Tumour necrosis factor receptor associated periodic syndrome (TRAPS)
- Hyperimmunoglobulin D syndrome (HIDS)/mevalonate kinase deficiency (MKD)
- Familial Mediterranean Fever (FMF)
- ILARIS should be given in combination with colchicine, if appropriate

#### Still's disease

ILARIS is indicated in patients who have responded inadequately to previous therapy with non-steroidal anti-inflammatory drugs (NSAIDs) and systemic corticosteroids, for the treatment of active Still's disease, including:

- Adult-onset Still's disease (AOSD)
- Systemic juvenile idiopathic arthritis (SJIA) in patients aged 2 years and older
- ILARIS can be given as monotherapy or in combination with methotrexate

#### Gouty arthritis

ILARIS is indicated for the symptomatic treatment of adult patients with frequent gouty arthritis attacks (at least 3 attacks in the previous 12 months) in whom non-steroidal anti-inflammatory drugs (NSAIDs) and colchicine are contraindicated, are not tolerated, or do not provide an adequate response, and in whom repeated courses of corticosteroids are not appropriate.

---

## **ILARIS safety profile in periodic fever syndromes (PFS)**

**FMF, HIDS/MKD, TRAPS and CAPS**

[Find out more](#)

## ILARIS safety profile in Still's disease

### AOSD and SJIA

[Find out more](#)

---

AOSD, adult-onset Still's disease; CAPS, cryopyrin-associated periodic syndromes; CINCA, chronic infantile neurological, cutaneous, articular syndrome; FCAS, familial cold autoinflammatory syndrome; FCU, familial cold urticaria; FMF, familial Mediterranean fever; HIDS, hyperimmunoglobulin D syndrome; MKD, mevalonate kinase deficiency; MWS, Muckle-Wells syndrome; NOMID, neonatal-onset multisystem inflammatory disease; NSAID, non-steroidal anti-inflammatory drug; SJIA, systemic juvenile idiopathic arthritis; TRAPS, tumour necrosis factor receptor-associated periodic syndrome.

### Reference

1. ILARIS® (canakinumab) Summary of Product Characteristics.



---

**Safety profile in PFS**

**Safety profile in PFS**

See more details

Hide details

UK | March 2025 | FA-11324189

Adverse events should be reported. Reporting forms and information can be found at [www.mhra.gov.uk/yellowcard](http://www.mhra.gov.uk/yellowcard). Adverse events should also be reported to Novartis online through the pharmacovigilance intake (PVI) tool at [www.novartis.com/report](http://www.novartis.com/report), or alternatively email [medinfo.uk@novartis.com](mailto:medinfo.uk@novartis.com) or call 01276 698370.

---

**Source URL:**

<https://www.pro.novartis.com/uk-en/medicines/immunology/ilaris/safety-profile>